



*Helping You Live Healthier Every Day*

FACTSHEET   FACTSHEET   FACTSHEET   FACTSHEET   FACTSHEET   FACTSHEET   FACTSHEET   FACTSHEET

OFFICE OF PUBLIC HEALTH • Center for Health Policy, Information & Promotion

Bioterrorism for the Public

# FAQ's - Hemorrhagic Fevers

---

## Facts about Viral Hemorrhagic Fever

Viral hemorrhagic fevers (VHFs) refer to a group of illnesses that are caused by several distinct families of viruses. In general, the term “viral hemorrhagic fever” is used to describe a severe multisystem syndrome (multisystem in that multiple organ systems in the body are affected). Characteristically, the overall vascular system is damaged, and the body’s ability to regulate itself is impaired. These symptoms are often accompanied by hemorrhage (bleeding); however, the bleeding is itself rarely life-threatening. While some types of hemorrhagic fever viruses can cause relatively mild illnesses, many of these viruses cause severe, life-threatening disease.

The Special Pathogens Branch (SPB) primarily works with hemorrhagic fever viruses that are classified as [biosafety level four](#) (BSL-4) pathogens. A list of these viruses appears in the [SPB disease information index](#). The Division of Vector-Borne Infectious Diseases, also in the National Center for Infectious Diseases, works with the non-BSL-4 viruses that cause two other hemorrhagic fevers, dengue hemorrhagic fever and yellow fever. Viruses causing hemorrhagic fever are initially transmitted to humans when the activities of infected reservoir hosts or vectors and humans overlap. The viruses carried in rodent reservoirs are transmitted when humans have contact with urine, fecal matter, saliva, or other body excretions from infected rodents. The viruses associated with arthropod vectors are spread most often when the vector mosquito or tick bites a human, or when a human crushes a tick. However, some of these vectors may spread virus to animals, livestock, for example. Humans then become infected when they care for or slaughter the animals.

Some viruses that cause hemorrhagic fever can spread from one person to another, once an initial person has become infected. [Ebola](#), [Marburg](#), [Lassa](#) and Crimean-Congo hemorrhagic fever viruses are examples. This type of secondary transmission of the virus can occur directly, through close contact with infected people or their body fluids. It can also occur indirectly, through contact with objects contaminated with infected body fluids. For example, contaminated syringes and needles have played an important role in spreading infection in outbreaks of Ebola hemorrhagic fever and Lassa fever.

Specific signs and symptoms vary by the type of VHF, but initial signs and symptoms often include marked fever, fatigue, dizziness, muscle aches, loss of strength, and exhaustion. Patients with severe cases of VHF often show signs of bleeding under the skin, in internal organs, or from body orifices like the mouth, eyes, or ears. However, although they may bleed from many sites around the body, patients rarely die because of blood loss. Severely ill patient cases may also show shock, nervous system malfunction, coma, delirium, and seizures. Some types of VHF are associated with renal (kidney) failure.

## Links

- [Viral Hemorrhagic Fevers](#)  
<http://www.cdc.gov/ncidod/dvrd/spb/mnpages/dispages/vhf.htm>
- [Viral Hemorrhagic Fevers Information](#)  
<http://www.bt.cdc.gov/agent/vhf/index.asp>